

Original article

On Fuzzy Subhypernear-Ring

Yasser AbouElwan*^{ID}, Sanaa Qabous, Samah Qadous

Department of Mathematics, Faculty of Science, University of Omar Almukhtar, Libya

ARTICLE INFO

Corresponding Email. yasser.abouelwan@omu.edu.ly

Received: 19-05-2024

Accepted: 26-06-2024

Published: 30-06-2024

Keywords. Near-Ring, Subnear-Ring, Level Set, Fuzzy Set, Hyper Near-Ring, Fuzzyhypernear-Ring.

Copyright: © 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution International License (CC BY 4.0). <http://creativecommons.org/licenses/by/4.0/>

Cite this article. AbouElwan Y, Qabous S, Qadous S. On Fuzzy Subhypernear-ring. Alq J Med App Sci. 2024;7(3):449-451. <https://doi.org/10.54361/ajmas.247303>

ABSTRACT

In this study, two main goals were achieved. In near-ring N , we have proven that, the level set is a subnear-ring of N if and only if the fuzzy set is the fuzzy subnear-ring of N . Similarly, in hyper near-ring R , it has been proven that, a fuzzy set is a fuzzy hypernear-ring if and only if level set is a subhypernear-ring of R . A direct proof method was used to reach these results, which will contribute to expanding the field of study on fuzzy near-ring and fuzzy hyper near-rings.

INTRODUCTION

The fuzzy sets on hyperstructures were first introduced by Zadeh in 1965 [1]. The fuzzy sets and algebraic hyperstructures have been considered by, Abou-Zaid [2], Davvaz [3] and others. Fuzzy hyperideals of hypernear-rings are a notion that Davvaz presented along with certain associated properties.

Fuzzy hyperideals of hypernear-rings are a notion that Davvaz presented along with certain associated properties. In this work, we will discuss relationships between fuzzy subset and level set on near-ring (hypernear-ring) respectively.

Near-ring

Definition 1.1.[6] A left near-ring is an algebraic structure $(N, +, \cdot)$ which satisfies the following axioms:

- (i) $(N, +)$ is a group (not necessarily abelian),
- (ii) with respect to the multiplication, (N, \cdot) is a semi-group,
- (iii) the multiplication is distributive with respect to the addition on the left side, i.e. $z \cdot (x + y) = z \cdot x + z \cdot y$ for all $x, y, z \in N$

Or right near-ring if satisfies the right distributive law.

$$(x + y) \cdot z = x \cdot z + y \cdot z, \text{ for all } x, y, z \in N.$$

The term "near-ring" will be used to refer to "left near-ring."

Example 1.2. $(\mathbb{Z}_8, +)$ is a group under '+' modulo 8.

Define ' \cdot ' on \mathbb{Z}_8 by $a \cdot b = a$ for all $a, b \in \mathbb{Z}_8$. Clearly $(\mathbb{Z}_8, +, \cdot)$ is a near-ring.

Definition 1.3.[4] A subgroup M of an near-ring N with $M \cdot M \subseteq M$ is called a **subnear-ring** of N , ($M \leq N$). A subgroup S of N with $N \cdot S \subseteq S$ is called a **normal subgroup** of N , ($S \trianglelefteq N$).

Hyper near-ring

Definition 2.1.[6] Let H be a nonempty set. A map $\circ : H \times H \rightarrow P^*(H)$ is called **hyper-operation**, $P^*(H)$ is the family of all nonvoid subsets of H .

Definition 2.2.[9] The triple $(R, +, \cdot)$ is a **hypernear-ring** if:

- I) $(R, +)$ satisfies the following axioms:
 - (1) $x + (y + z) = (x + y) + z$, for any $x, y, z \in R$.

- (2) $\exists 0 \in R$ s.t. for any $x \in R$, $x + 0 = 0 + x = x$.
 (3) for any $x \in R$, there exists a unique element $-x \in R$, such that

$$0 \in x + (-x) \cap -x + x.$$

 (4) for any $x, y, z \in R$, $z \in x + y$ implies that $x \in z + (-y)$, $y \in -x + z$.
 II) (R, \cdot) is a semi-group endowed with a two-sided absorbing element 0, i.e. for any $x \in R$, $x \cdot 0 = 0 \cdot x = 0$.
 III) The operation ' \cdot ' is distributive with respect to the hyperoperation '+' from the left side: For many $x, y, z \in R$,
 $x \cdot (y + z) = x \cdot y + x \cdot z$,

Definition 2.3.[7] Let $(R, +, \cdot)$ be a hypernear-ring. A non-empty subset H of R is called a subhypernear-ring if

- (1) $(H, +)$ is a subhypergroup of $(R, +)$, i.e., $a, b \in H$ implies $a + b \subseteq H$, and $a \in H$ implies $-a \in H$,
 (2) $ab \in H$, for all $a, b \in H$.

Example 2.4.[9] Let $R = \{0, a, b, c\}$ be a set with a hyperoperation "+" and a binary operation " \cdot " as follows:

Table 2.1

+	0	a	b	c
0	{0}	{a}	{b}	{c}
a	{a}	{0,a}	{b}	{c}
b	{b}	{b}	{0,a,c}	{b,c}
c	{c}	{c}	{b,c}	{0,a,b}

Table 2.2

\cdot	0	a	b	c
0	0	a	b	c
a	0	a	b	c
b	0	a	b	c
c	0	a	b	c

Then $(R, +, \cdot)$ is a hypernear-ring, and $\{0\}$, $\{0, a\}$, and R are subhypernear-rings of R .

Fuzzy structure

Definition 3.1.[8] A *fuzzy subset* of X is a function $\mu: X \rightarrow [0, 1]$. The set of all fuzzy subsets of X is called the fuzzy power set of X and is denoted by $FP(X)$. For $t \in [0, 1]$, define μ_t as follows:

$$\mu_t = \{x \mid x \in X, \mu(x) \geq t\}. \mu_t \text{ is called the } t\text{-level set of } \mu.$$

Example 3.2. In Example 2.4. $R = \{0, a, b, c\}$, define a fuzzy subset $\mu: R \rightarrow [0, 1]$ by: $\mu(0) = 1$, $\mu(a) = 0.7$, $\mu(b) = \mu(c) = 0.3$.

Note that $\mu_{0.45} = \{x \in U \mid \mu(x) \geq 0.45\} = \{0, a\}$ and $\mu_0 = R$.

Definition 3.3.[7] Let N be a near-ring and μ be a fuzzy subset of N . We say μ a *fuzzy subnear-ring of N* if for all $x, y \in N$,

- (1) $\mu(x - y) \geq \min \{\mu(x), \mu(y)\}$,
 (2) $\mu(xy) \geq \min \{\mu(x), \mu(y)\}$.

Theorem 3.4. Let N be a near-ring and μ be a fuzzy subset of N . Then the level subset $\mu_t (\neq \emptyset)$ is a subnear-ring of N for all $t \in (0, 1]$ if and only if μ is a fuzzy subnear-ring of N .

Proof. Let μ_t is a subnear-ring of N .

Let $x, y \in N$ and putting $t_0 = \min \{\mu(x), \mu(y)\}$ then $x, y \in \mu_{t_0}$

$\Rightarrow \mu(x) \geq t_0, \mu(y) \geq t_0$, and since μ_t is a subnear-ring of N , hence $xy \in \mu_{t_0}$, and so $\mu(xy) \geq t_0 = \min \{\mu(x), \mu(y)\}$, also, $x - y \in \mu_{t_0}$ so

$\mu(x - y) \geq t_0 = \min \{\mu(x), \mu(y)\}$, therefore μ is fuzzy subnear-ring.

Conversly, Let μ is fuzzy subnear-ring and $t \in (0, 1]$, $\mu_t (\neq \emptyset)$

Let $x, y \in \mu_t \Rightarrow \mu(x) \geq t, \mu(y) \geq t$, then $\min \{\mu(x), \mu(y)\} \geq t$.

And $\mu(xy) \geq \min \{\mu(x), \mu(y)\} \geq t$ (since μ is fuzzy subnear-ring)

Therefore $xy \in \mu_t$, hence $\mu_t \cdot \mu_t \subseteq \mu_t$. Thus, μ_t is a subnear-ring of N . ■

Definition 3.5.[5] Let $(R, +, \cdot)$ be a hypernear-ring. Then we call a fuzzy set μ of R a *fuzzy subhypernear-ring of R* if it satisfies the following inequalities:

- (1a) $\min \{\mu(x), \mu(y)\} \leq \inf_{z \in x+y} \mu(z)$ for all $x, y \in R$,
 (1b) $\mu(x) \leq \mu(-x)$ for all $x \in R$,
 (2) $\min \{\mu(x), \mu(y)\} \leq \mu(xy)$ for all $x, y \in R$.

Theorem 3.6. A fuzzy set μ of R is a fuzzy subhypernear-ring of R if and only if for any $t \in [0, 1]$, $\mu_t (\neq \emptyset)$ is a subhypernear-ring of R .

Proof. Let μ is a fuzzy sub-hypernear-ring and $t \in [0, 1]$, $\mu_t \neq \emptyset$

If $x, y \in \mu_t \Rightarrow \mu(x) \geq t, \Rightarrow \mu(y) \geq t$

Hence $\inf \mu(z) \geq \min\{\mu(x), \mu(y)\} \geq t$ where $z \in x + y$

Therefore, for all $z \in x + y$ we have $z \in \mu_t$ and so $x + y \subseteq \mu_t$

Thus μ_t is sub-hypernear-ring of R

Conversely, let μ_t is sub-hypernear-ring of R

Let $x, y \in R$ and putting $t_0 = \min\{\mu(x), \mu(y)\}$ then $x, y \in \mu_{t_0}$

Since μ_t is sub-hypernear-ring of R

$\therefore x + y \subseteq \mu_t \Rightarrow$ for any $z \in x + y, z \in \mu_{t_0}$ which implies that

$\inf \mu(z) \geq t_0 = \min\{\mu(x), \mu(y)\}$ where $z \in x + y$

and since μ_t is a sub-hypernear-ring of R, hence $xy \in \mu_t$

$$\mu(xy) \geq t_0 = \min\{\mu(x), \mu(y)\}$$

Thus μ of R is a fuzzy subhypernear-ring of R. ■

Example 3.7. In Example 2.4., and Example 3.2.

$$\mu_t = \begin{cases} R, & t \in (0, 0.3] \\ \{0, a\}, & t \in (0.3, 0.7] \\ \{0\}, & t \in (0.7, 1] \end{cases}$$

Clearly, all μ_t are subhypernear-ring of R, and μ is a fuzzy subhypernear-ring of R.

CONCLUSION

In this paper, the following has been proven that the level set of near-ring is a subnear-ring if and only if the fuzzy set is the fuzzy subnear-ring. A fuzzy set of hyper near-ring is a fuzzy hypernear-ring if and only if level set is a sub-hypernear-ring.

REFERENCES

1. Abou-Zaid S. On fuzzy sub-near-rings and ideals, Fuzzy Sets Syst. 1991;44:139-146.
2. Davvaz B. Fuzzy Algebraic Hyperstructures, Springer, 2005. DOI 10.1007/978-3-319-14762-8.
3. Stayanarayana B, Preasad K. Near rings, fuzzy ideals, and graph theory. 2013. CRC Press.
4. Davvaz B, Zhan J, Kim K. Fuzzy Γ -hypernear-rings. Computers and Mathematics with Applications. 2010;59(8):2846-2853.
5. AbouElwan Y, Alderawe A. On Homomorphisms and Congruences of Canonical Hypergroups, Global Libyan Journal. 2023;68:1-8.
6. Davvaz B. $(\epsilon, \epsilon \vee q)$ – fuzzy subnear-rings and ideals, Soft Comput. 2006;10:206-211.
7. Mordeson J, Bhutani K, Rosenfeld A. Fuzzy Group Theory, Springer-Verlag Berlin Heidelberg. 2005
8. AbouElwan Y, Fanoush M, Elmabrouk T. On Hyper Ideals of Γ – hypernear-ring. Global Libyan Journal. 2020;4:25-45.

حول قريب الحلقة الفوقية الجزئية الضبابية

ياسر أحمد أبوعلوان، سناء حمدي قادوس، سماح حمدي قادوس
قسم الرياضيات، كلية العلوم، جامعة عمر المختار، ليبيا

المستخلص:

في هذه الدراسة، تم تحقيق هدفين، في قريب الحلقة، تم إثبات أن، المجموعة المنسوبة تكون قريب الحلقة الجزئية لقريب الحلقة اذا واذا فقط كانت المجموعة الضبابية هي قريب الحلقة الضبابية. بالمثل في قريب الحلقة الضبابية تم إثبات أن، المجموعة الضبابية تكون قريب الحلقة الضبابية اذا واذا فقط كانت المجموعة المنسوبة هي قريب الحلقة الفوقية الجزئية لقريب الحلقة الفوقية. تم استخدام أسلوب البرهان المباشر للوصول لهذه النتائج، التي ستسهم في توسيع مجال الدراسة حول قريب الحلقات الضبابية والفوقية الضبابية.

الكلمات الدالة: قريب الحلقة، قريب الحلقة الجزئية، المجموعة المنسوبة، المجموعة الضبابية، قريب الحلقة الفوقية، قريب الحلقة الفائقة الضبابية.