

Original article

Synthesis of Cyclopentanone and Cyclohexanone Derivatives

Naowara Alarafi^{*}, Abdullah Gheath, Abdalhamid Alarafi

Department of Chemistry, Faculty of Science, Benghazi University, Benghazi, Libya

Corresponding Email. Nourapro@gmail.com

Abstract

A series of compounds (I–XIII) were synthesized by reacting ethyl 2-oxocyclopentane-1-carboxylate (1) or ethyl 2-oxocyclohexane-1-carboxylate (2) with various diamines, including ethane-1,2-diamine, benzene-1,4-diamine, hydrazine hydrate, benzidine, N¹, N³-bis(2-aminoethyl) malonamide, 3-hydroxyaniline, malonohydrazide, and piperazine. Most derivatives (I, II, III, VI, VII, VIII, X, XIII) adopted enamine structures via a 2:1 reaction ratio, whereas compounds IX and XI exhibited distinct structural frameworks due to a 1:1 reaction pathway. Notably, the reaction of β -keto ester 1 with hydrazine hydrate yielded enamine III as a minor product and pyrazolone IV as the major derivative, while the analogous reaction with compound (2) exclusively produced pyrazolone V. Additionally, the reaction of compound (2) with malonohydrazide afforded an azine derivative (XII). All syntheses proceeded under mild conditions with moderate yields, and the structures were characterized using melting point, TLC, IR, MS, ¹H-NMR (with D₂O exchange), ¹³C-NMR, and APT analyses. Preliminary screening revealed antimicrobial activity in selected derivatives.

Keywords. Diamines, Cyclopentanone Derivatives, Secondary Amines, Neat Synthesis.

Introduction

The structural analogs, enamines and enols, are produced when secondary amines react with carbonyl compounds. Pyrrolidine, piperidine, and morpholine are common secondary amines that are used to create enamines [1]. Enamines are substances with a lot of electrons and interesting redox characteristics. Here, several secondary enamines that were condensed from primary amine and β -keto carbonyls were created, and cyclic voltammetry was used to methodically examine their electrochemical oxidation characteristics [2]. The first general synthesis of enamines was published by Mannich nine years before [3]. Wittig created the term "enamine" in 1927 to highlight the similarity between this family of chemicals and enols [4], although the first known instances of enamine chemistry were in 1884 [5-7]. It wasn't until 1954 that Stork's groundbreaking research revealed the synthetic potential of enamine-electrolyte interaction [8-11].

Enamine chemistry has been thoroughly examined [12]. Before the early 1970s, enamines held a special position as noncharged enolate equivalents; since then, they have shared this status with silylated enol ethers [13-20]. Their reactivity with acceptor-activated aryl halides [21], electron-deficient dienes π 4 cycloaddition partners [22-24], and Michael acceptors [25-26]. Controllable and responsive systems may be created by constitutional exchange utilizing dynamic interactions and reactions of either a covalent or non-covalent nature [27-44]. Enamine compounds are widely recognized for their diverse biological activities, attributed to their unique structural motif—a nitrogen-containing group conjugated with a carbon-carbon double bond. This configuration enables enamines to serve as versatile intermediates in organic synthesis while facilitating strong interactions with biological targets. Extensive studies have demonstrated that enamine derivatives exhibit a broad spectrum of pharmacological properties, including antibacterial, antifungal, antiviral, anti-inflammatory, and anticancer activities [45]. Given their pharmacological relevance, this study focuses on the synthesis of new enamine derivatives using an innovative, facile, and high-yield preparation method. The approach aims to expand the library of bioactive enamines for further exploration of their therapeutic potential.

Materials and Methods

Materials

All chemicals were purchased from SigmaAldrich (St. Louis, MO, USA) and were used as received. Reactions were monitored on TLC (Chloroform /pet.methanol). All spectroscopic analysis of prepared compounds was conducted in the National Research Centre at Giza. ¹H NMR spectra were carried out on a Bruker 400 MHz with chemical shift (δ) expressed in ppm downfield from tetramethyl silane as an internal standard (δ TMS = 0) using CDCl₃ and DMSO-d₆ as solvents. The multiplicity of the signal is as follows: s (Singlet), d (Doublet), t (Triplet), q (Quartet), m (Multiplet). ¹³C-NMR was measured on a Bruker 100 MHz with internal reference TMS δ = 0. Infrared spectra were recorded on a Perkin Elmer 2000 FT-IR system spectrometer, where the positions of absorptions have been expressed in wave number units (cm⁻¹). Melting points (m.p) of the synthesized compounds were measured in capillary tubes using a Stuart scientific apparatus and are uncorrected.

Synthetic procedure**Synthesis of enamines (I – VI)**

Two equivalents of either ethyl 2-oxocyclopentane-1-carboxylate or ethyl 2-oxocyclohexane-1-carboxylate were added to one equivalent of the amine. the mixture was stirred at room temperature under solvent-free conditions. The resulting precipitate was collected by filtration, washed with ethanol, and recrystallized from ethanol.

Synthesis of enamines (VII-XIII)

Two equivalents of ethyl-2-oxocyclopentane-1-carboxylate or ethyl 2-oxocyclohexane-1-carboxylate were mixed with one equivalent of the amine, and the mixture was refluxed in ethanol until precipitation was observed. The formed precipitate was then filtered, washed with ethanol, and recrystallized from ethanol.

Diethyl 2,2'-(ethane-1,2-diylbis(azanediyl)) bis (cyclopent-1-ene-1-carboxylate) (I): 36% yield as an off-white solid. m.p. 108-110 °C. IR (KBr) ν_{max} : 1591.53 cm^{-1} (C=C), 1644.40 cm^{-1} (ester CO), and 2945.33 cm^{-1} (CH aliphatic), 3317.45 cm^{-1} (NH amine). ^1H NMR (CDCl_3): δ = 7.43 (2H, s, NH), 4.12 (4H, q, CH_2CH_3), 3.29 (4H, s, $\text{CH}_2\text{-NH}$), 2.47-2.41 (8H, dd, CH_2), 1.79 – 1.74 (4H, m, CH_2), 1.26 (6H, t, CH_2CH_3). ^{13}C NMR (CDCl_3): δ = 168.45 (CO), 164.16 (C=C-NH), 93.82, 58.49, 46.00, 31.94, 29.07, 20.88, 14.68. MS EI m/z: M^+ . 336, 168, 291, 245, 122, 94.

Diethyl 2,2'-(ethane-1,2-diylbis(azanediyl)) bis (cyclohexene-1-carboxylate) (II): 87% yield as a white solid. m.p. 70-72 °C. IR (KBr) ν_{max} : 1582.55 cm^{-1} (C=C), 1637.54 cm^{-1} (ester CO), 2973.32 cm^{-1} (CH aliphatic), 3252.81 cm^{-1} (NH amine). ^1H NMR (CDCl_3) δ = 8.99 (2H, s, NH), 4.12 (4H, q, CH_2CH_3), 3.29 (4H, dd, $\text{CH}_2\text{-NH}$), 2.26 (8H, m, CH_2), 1.62 (8H, m, CH_2), 1.26 (6H, t, $\text{CH}_3\text{-CH}_2$). ^{13}C NMR (CDCl_3): δ = 170.86 (CO), 159.03 (C=C-NH), 90.60, 58.66, 43.09, 26.24, 23.81, 22.59, 22.26, 14.62. MS EI m/z: M^+ . 364, 320, 292, 264, 198, 183, 158, 155.

Ethyl(Z)-2-(2-(2-(ethoxycarbonyl) cyclopentylidene)hydrazineyl)cyclopent-1-ene-1-carboxylate (III): 14% yield as a yellow solid. m.p. 71-73°C. IR (KBr) ν_{max} : 1604.55 cm^{-1} (C=C), 1647.36 and 1732.91 cm^{-1} (ester CO), and 2996 cm^{-1} (CH aliphatic), 3259.78 cm^{-1} (NH amine), ^1H NMR (CDCl_3): δ = 10 (1H, s, NH), 4.14-4.26 (4H, m, CH_2CH_3), 3.49 (1H, t, CH), 1.83-3.15 (12H, m, CH_2), 1.26-1.30 (6H, m, CH_2CH_3), ^{13}C NMR (CDCl_3): δ = 173.54 (CO), 169.42 (CO), 161.20 (C=N), 155.28 (N-C=C), 96.15 (C=C), 61.97, 61.36, 38.06, 32.15, 27.38, 25.95, 25.91, 23.22, 21.44, 14.67. MS m/z: 308.09. MS EI m/z: M^+ . 308, 263, 216, 188, 160, 108, 80.

1,4,5,6-tetrahydrocyclopenta[c]pyrazol-3(2H)-one (IV): 82% yield as a yellow solid. m.p. 243- 245°C. IR (KBr) ν_{max} : 3400 – 2500 cm^{-1} (OH), 3000 cm^{-1} (CH aliphatic), 1590 cm^{-1} (C=N), and 1529 cm^{-1} (C=C). ^1H NMR (DMSO-d_6): δ = 10.51 (2H, s, br., NH), 2.46 (2H, m, CH_2), 2.39-2.33 (4H, m, CH_2), ^{13}C NMR (DMSO-d_6): δ = 154.50 (CO), 153.50 (C=C-NH), 107.50 (C=C-CO), 30.47, 24.48, 22.62.

1,2,4,5,6,7-hexahydro-3H-indazol-3-one (V): 48% yield as a white solid. m.p. 268-270 °C. IR (KBr) ν_{max} : 3400 – 2500 cm^{-1} (OH), 3000 cm^{-1} (CH aliphatic), 1611 cm^{-1} (NH), 1560 cm^{-1} (C=N) and 1540 cm^{-1} (C=C). ^1H NMR (DMSO-d_6): δ = 10.26 (2H, s, br., NH), 2.5 (2H, t, CH_2), 2.41 (2H, t, CH_2), 1.62 (4H, m, CH_2), ^{13}C NMR (DMSO-d_6): δ = 158.95 (CO), 140.30 (C=C-NH), 98.94 (C=C-CO), 23.32, 22.75, 21.74, 19.35 MS EI m/z: M^+ . 138, 110, 81

Diethyl 2,2'-(piperazine-1,4-diyl) bis(cyclopent-1-ene-1-carboxylate) (VI): 85% yield, an off-white solid. m.p. 112-113 °C. IR (KBr) ν_{max} : 1666.82 cm^{-1} (C=C), 1742.13 cm^{-1} (ester CO), and 2981.90 cm^{-1} (CH aliphatic). ^1H NMR (CDCl_3): δ = 4.13 (4H, q, CH_2CH_3), 3.48 (6H, m, CH_2), 2.64 (8H, m, CH_2), 1.79 (4H, m, CH_2) 1.26 (6H, t, CH_3). ^{13}C NMR (CDCl_3): δ = 165.88 (CO), 161.79 (C=C-N), 103.09 (C=C-CO), 97.63 (C=C-CO), 61.31, 60.48, 49.88, 49.09, 35.25, 32.38, 20.93, 20.18, 14.64, 14.14. MS EI m/z: M^+ . 362, 333, 289, 243, 224, 194, 135, 122, 110.

Diethyl 2,2'-(1,4-phenylenebis(azanediyl)) bis(cyclopent-1-ene-1-carboxylate) (VII): 97% yield as a purple solid. m.p. 164-165°C. IR (KBr) ν_{max} : 1601.08 cm^{-1} (C=C), 1653.82 cm^{-1} (ester CO), 2855.67 cm^{-1} (CH aliphatic), 3055.82 (=CH), 3295.97 cm^{-1} (NH amine), ^1H NMR (CDCl_3): δ 9.53 (2H, s, NH), 6.97 (4H, s, Ar-CH), 4.23 (4H, q, CH_2CH_3), 2.73 (4H, t, CH_2), 2.55-2.76 (4H, t, CH_2), 1.87 (4H, q, CH_2), 1.31 (6H, t, CH_2CH_3), ^{13}C NMR (CDCl_3): δ = 168.49. (CO), 160.57 (C=C-NH), 136.36, 121.90, 97.28, 58.94, 33.51, 28.79, 21.73, 14.68. MS EI m/z: M^+ . 384, 338, 292, 264, 237, 184, 156.

Diethyl 2,2'-(1,4-phenylenebis(azanediyl))bis(cyclohex-1-ene-1-carboxylate) (VIII): 36% yield as an off-gray color. m.p. 153- 154°C. IR (KBr) ν_{max} : 1589.54 cm^{-1} (C=C), 1643.36 cm^{-1} (ester CO), 2978.18 cm^{-1} (CH aliphatic), 3152.51 cm^{-1} (=CH), 3207.53 cm^{-1} (NH amine). ^1H NMR (CDCl_3): δ = 10.70 (2H, s, NH), 6.98

(4H, s, Ar-CH), 4.22 (4H, q, CH₂CH₃), 2.37 (8H, t, CH₂), 1.62 (8H, m, CH₂), 1.29 (6H, t, CH₂CH₃). ¹³C NMR (CDCl₃): δ = 170.84 (CO), 156.58 (C=C-NH), 136.23, 125.38, 93.04, 57.22, 29.96, 27.09, 23.85, 22.32, 14.59. MS EI m/z: M⁺. 412, 364, 319, 136, 108, 79.

Ethyl 2-((4'-amino-[1,1'-biphenyl]-4-yl) amino) cyclopent-1-ene-1-carboxylate (IX): 12% yield as a yellow solid. m.p. 169-170°C. ¹H NMR (CDCl₃): δ = 9.70 (1H, s, NH) 7.53-7.48 (8H, m, Ar-CH), 4.25 (2H, q, CH₂CH₃), 3.60 (2H, s, NH₂), 2.89 (2H, t, CH₂), 2.06 (2H, t, CH₂) 1.93(2H, quintet, CH₂), 1.34 (3H, t, CH₂CH₃). ¹³C NMR (CDCl₃): δ = 168.53, 160.47, 139.80, 135.03, 127.58, 126.98, 120.79, 115.46, 98.09, 59.03, 33.77, 28.74, 21.84, 14.68. MS EI m/z: M⁺. 322, 276, 250, 184, 167, 138, 123, 77.

Diethyl 2,2'-([1,1'-biphenyl]-4,4'-diylbis(azanediyl)) bis(cyclopent-1-ene-1-carboxylate)

(X): 18% yield as a yellow solid. m.p. 154-160°C. ¹H NMR (CDCl₃): δ = 9.69 (1H, s, NH) 6.84-7.50 (8H, m, Ar-CH), 4.26 (4H, q, CH₂CH₃), 2.87 (4H, t, CH₂), 2.60 (4H, t, CH₂), 1.92 (4H, quintet, CH₂), 1.34 (6H, t, CH₂CH₃) [enamine]. ¹³C NMR (CDCl₃): δ = 173.55, 165.40, 137.85, 129.96, 125.33, 118.83, 54.64, 37.48, 33.30, 29.49, 21.82, 19.82[imine].

Ethyl 2-((4'-amino-[1,1'-biphenyl]-4-yl) amino) cyclohex-1-ene-1-carboxylate (XI): 43% yield as a yellow solid. m.p. 162-164°C. IR (KBr) u_{max}: 1578.29 cm⁻¹ (C=C), 1631.60 cm⁻¹ (ester CO), 2980.96 cm⁻¹ (CH aliphatic), 3036.16 (=CH) 3223.09 cm⁻¹ (NH amine), 3350.92, 3453.78 (NH₂). ¹H NMR (CDCl₃): δ = 10.87 (1H, d, NH) 6.75-7.52 (8H, m, Ar-CH), 4.24 (2H, q, CH₂CH₃), 3.70 (2H, s.br., NH₂), 2.40 -1.62 (8H, m, CH₂), 1.32 (3H, t, CH₂CH₃). MS EI m/z: M⁺. 336, 290, 234, 184, 167, 145, 91.

1,2,3,4,6,7,9,10,11,12,13,14-dodecahydrodibenzo [c, i][1,2,6,7]tetrazecine-5,8-dione (XII): 24% yield as an off-white solid. m.p. 280-282°C. ¹H NMR (DMSO-d₆): δ = 10.27 (4H, s.br., NH), 2.50-2.41 (4H, t, CH₂), 2.24-2.21 (4H, t, CH₂), 1.64-1.63 (8H, m, CH₂). ¹³C NMR (DMSO-d₆): δ = 159.04 (CO), 140.54 (C=C-NH), 99.03 (C=C-C=O), 23.30, 22.72, 21.74, 19.32.

Ethyl (Z)-2-((2-(3-((2-(ethoxycarbonyl) cyclopentylidene) amino) ethyl) amino)-3-oxopropanamido) ethyl) amino) cyclopent-1-ene-1-carboxylate (XIII): 40% yield as a white solid m.p. 85-88°C. IR (KBr) u_{max}: 1585.80 cm⁻¹ (C=C), 1642.43 cm⁻¹ (ester CO), 2945.91 cm⁻¹ (CH aliphatic), 3316.82 cm⁻¹ (NH amine). ¹H NMR (DMSO-d₆): δ = 8.161 (1H, s, NH), 7.452 (2H, s, NH) 4.031 (4H, q, CH₂CH₃), 3.02-3.65 (15H, m, CH₂), 2.35 -2.54(8H, m, CH₂), 1.18 (6H, t, CH₂CH₃) ¹³C NMR (DMSO-d₆): δ = 179.25, 167.62, 167.50, 165.24, 91.83, 58.12, 45.15, 43.70, 40.54, 39.70, 39.29, 20.87, 15.18.

Results and Discussion


This paper reviews the chemistry of enamine and imine derivatives, with a focus on the synthesis of enamines via the reaction of cyclic β-ketoesters with primary or secondary amines. In these reactions, amines act as nucleophiles, while cyclic β-ketoesters function as electrophiles, undergoing nucleophilic attack to yield a novel series of enamine-containing compounds (I-XIII). In this study, ethyl 2-oxocyclopentane-1-carboxylate (1) and ethyl 2-oxocyclohexane-1-carboxylate (2) were reacted with various amines—including ethane-1,2-diamine, benzidine, benzene-1,4-diamine, piperazine, hydrazine hydrate, N¹,N³-bis(2-aminoethyl)malonamide, and malonohydrazide—under neat conditions at room temperature and under reflux in ethanol.

Ethyl 2-oxocyclopentane-1-carboxylate (1) was reacted with a series of diamines, including ethane-1,2-diamine, hydrazine hydrate, piperazine, 1,4-phenylenediamine, benzidine, and N¹, N³-bis(2-aminoethyl) malonamide, yielding compounds I, III, IV, VI, VII, IX, X, and XIII, respectively as shown in scheme (1).

In a parallel study, ethyl 2-oxocyclohexane-1-carboxylate (2) was treated with the same set of diamines namely ethane-1,2-diamine, hydrazine hydrate, 1,4-phenylenediamine, benzidine, and malonohydrazide resulting in the synthesis of compounds II, V, VIII, XI, and XII, as shown in scheme (2). These reactions demonstrate the versatility of β-keto esters in forming diverse heterocyclic and condensed structures upon condensation with different diamines.




Compounds I, II, III, VI, VII, VIII, X, and XIII displayed nearly identical structural frameworks, with the reaction proceeding in a 2:1 stoichiometric ratio that is, two molecules of β -keto esters (1) or (2) reacted with one molecule of the diamine. In contrast, an analysis of the reaction schemes depicted in Figures 1 and 2 revealed that compounds IX and XI were formed via a 1:1 ratio reaction between β -keto esters (1) or (2) and benzidine. The proposed mechanism below scheme (3) outlines the reaction pathway of β -keto esters (1) or (2) with these diamines.




Scheme 3. The proposed mechanism of synthesis of compounds (I, II, III, VI, VII, VIII, X, XIII)

The reaction of ethyl 2-oxocyclopentanecarboxylate (1) with hydrazine hydrate proceeded via competing pathways, yielding two distinct products. A pyrazolone derivative (IV) was obtained as the major product (85% yield), while an enamine derivative (III) was isolated as the minor product (14% yield). In contrast, under identical conditions, ethyl 2-oxocyclohexane-1-carboxylate (2) reacted with hydrazine hydrate to exclusively form product V. A proposed mechanistic scheme (4) delineates the formation of IV and V, underscoring the divergent reactivity between the five- and six-membered ring systems.



Scheme 4: The proposed mechanism of synthesis of compounds (IV and V)

Additionally, the synthesis of a novel compound (XII) was achieved through the condensation of ethyl 2-oxocyclohexane-1-carboxylate (1) with malonohydrazide under reflux conditions. This transformation proceeded via the construction of a tetrazecine ring, as depicted in the mechanistic pathway (5). The observed product distribution suggests that the ring size significantly influences the reaction pathway, with the five-membered substrate favoring a competitive mechanism, while the six-membered analogue follows a more selective route. Further studies could explore the underlying steric and electronic factors governing these transformations.



Scheme 5. The proposed mechanism of synthesis of compound (XII)

The successful preparation of these compounds was confirmed using various analytical techniques, including infrared (IR) spectroscopy and nuclear magnetic resonance (NMR) spectroscopy (^1H -NMR and ^{13}C -NMR). The IR spectra of these compounds exhibited characteristic absorption bands corresponding to the ester carbonyl group ($\text{C}=\text{O}$) and the secondary amine ($\text{N}-\text{H}$) functionality. Additionally, the ^1H -NMR spectra displayed signals consistent with the secondary amine proton and the ethyl group of the ester moiety. A notable distinction was observed in the ^1H -NMR spectra of compounds (IV) and (V). Specifically, the spectra of (IV) and (V) lacked signals attributable to the ethoxy group, indicating nucleophilic attack on the ester moiety and subsequent formation of the pyrazolone rings. In contrast, the ^1H -NMR spectra of (IX) and (XI) exhibited a triplet at 1.34, 1.32 ppm and a quartet at 4.25, 4.24 ppm, respectively, confirming the retention of the ethoxy groups. Furthermore, a broad singlet at 3.60 ppm and 3.70 ppm, assigned to the amine protons (NH_2), suggested a 1:1 reaction stoichiometry with complete conversion. Meanwhile, the ^1H NMR spectrum of compound (XII) was also confirmed by the absence of the ethoxy group and the appearance of a broad signal representing four protons at 10.27 ppm, corresponding to the $\text{N}-\text{H}$ protons in the azine ring. Furthermore, ^{13}C -NMR analysis verified the presence of the ester carbonyl carbon in selected compounds, while it was absent in others with a peak corresponding to the $\text{C}=\text{C}-\text{NH}$ fragment. Additionally, the number of carbon signals in each compound's ^{13}C -NMR spectrum matched the expected number of carbons, further verifying the successful synthesis of these compounds. These combined spectroscopic analyses provided conclusive evidence for the successful preparation of the target compounds.

Conclusion

In this study, a series of novel enamine derivatives were successfully synthesized through the reaction of ethyl 2-oxocyclopentane-1-carboxylate (1) or ethyl 2-oxocyclohexane-1-carboxylate (2) with various diamines under mild conditions, yielding the target products with high efficiency. Structural elucidation via spectroscopic techniques (IR, ^1H NMR, ^{13}C NMR, and MS) confirmed the formation of the desired enamine derivatives. The reactivity of the substrates was found to be highly dependent on ring size and diamine structure. Compound (2), featuring a six-membered ring, predominantly formed single products with ethane-1,2-diamine, 1,4-phenylenediamine, and benzidine, adhering to 1:2 or 1:1 stoichiometries, respectively. In contrast, reactions with hydrazine hydrate and malonohydrazide afforded cyclic products. Compound (1), with its more strained five-membered ring, exhibited enhanced reactivity, leading to competitive pathways in some cases. For instance, reactions with benzidine yielded a mixture of 1:1 and 1:2 adducts, while hydrazine hydrate produced both pyrazolone and enamine derivatives. Other diamines, however, selectively afforded single products via 1:2 addition. The observed differences in reactivity and product distribution highlight the influence of ring strain and steric factors on the reaction pathways. The higher reactivity of the cyclopentane derivative (1) compared to the cyclohexane analogue (2) underscores the role of ring tension in governing mono- versus bis-addition and the preference for cyclic versus acyclic structures. These findings provide valuable insights into the design and synthesis of enamine-based derivatives with tailored structural and functional properties.

Conflict of interest. Nil

References

- Ouellette RJ, Rawn JD. Organic chemistry study guide: Key concepts, problems, and solutions. Amsterdam: Elsevier; 2015.
- Li Y, Wang D, Zhang L, Luo S. Redox property of enamines. J Org Chem. 2019;84(18):12071-90.
- Mannich C, Davidsen H. Über einfache Enamine mit tertiär gebundenem Stickstoff. Ber Dtsch Chem Ges (A B Ser). 1936;69(9):2106-12.
- Wittig G, Blumenthal H. Über die Einwirkung von Ammoniak und Ammoniak-Derivaten auf o-Acetylaceto-phenole. Ber Dtsch Chem Ges (A B Ser). 1927;60(5):1085-94.
- Collie JN. Ueber die Einwirkung des Ammoniaks auf Acetessigester. Justus Liebigs Ann Chem. 1884;226(3):294-322.
- Benary E. Über die Acylierung des β -Amino-crotonsäureesters und verwandter Verbindungen. Ber Dtsch Chem Ges. 1909;42(3):3912-25.
- Robinson R. The C-alkylation of certain derivatives of β -aminocrotonic acid. J Chem Soc. 1916;109:762-8.
- Stork G, Terrell R, Szmuszkowicz J. A new synthesis of 2-alkyl and 2-acyl ketones. J Am Chem Soc. 1954;76(7):2029-30.
- Stork G, Landesman HK. A new alkylation of carbonyl compounds. II. J Am Chem Soc. 1956;78(19):5128-9.
- Stork G, Landesman HK. A new ring enlargement sequence. J Am Chem Soc. 1956;78(19):5129-30.
- Stork G, Brizzolara A, Landesman H, Szmuszkowicz J, Terrell R. The enamine alkylation and acylation of carbonyl compounds. J Am Chem Soc. 1963;85(2):207-22.
- Huang ZT, Wang MX. The Chemistry of Enamines. New York: Wiley; 1994. p. 1303.
- Brownbridge P. Silyl enol ethers in Synthesis—part I. Synthesis. 1983;1983(1):1-28.
- Brownbridge P. Silyl enol ethers in synthesis—part II. Synthesis. 1983;1983(2):85-104.
- ReiSig HU. Siliciumverbindungen in der organischen Synthese. Chem Unserer Zeit. 1984;18(2):46-54.
- Reetz MT. Lewis-Säure-bewirkte α -Alkylierung von Carbonylverbindungen. Angew Chem. 1982;94(2):97-109.

17. Reetz MT. Lewis Acid Induced α -Alkylation of Carbonyl Compounds. *Angew Chem Int Ed Engl.* 1982;21(2):96-108.
18. Kaur P, Khatik GL, Nayak SK. A review on advances in organoborane chemistry: versatile tool in asymmetric synthesis. *Curr Org Synth.* 2017;14(5):665-82.
19. Reißig HU. Donor-acceptor-substituted cyclopropanes: Versatile building blocks in organic synthesis. In: de Meijere A, editor. *Small ring compounds in organic synthesis III.* Berlin: Springer; 1988. p. 73-135.
20. Mahrwald R. Diastereoselection in Lewis-acid-mediated aldol additions. *Chem Rev.* 1999;99(5):1095-120.
21. Kuehne ME. The arylation of enamines. *J Am Chem Soc.* 1962;84(5):837-47.
22. Shono T, Matsumura Y, Inoue K, Ohmizu H, Kashimura S. Electroorganic chemistry. 62. Reaction of iminium ion with nucleophile: A versatile synthesis of tetrahydroquinolines and julolidines. *J Am Chem Soc.* 1982;104(21):5753-7.
23. Murahashi SI, Naotao T, Nakato T. Convenient Method for the Construction of Quinoline Skeletons from N-Methylarylamines via N-(t-Butyldioxymethyl) arylamines. *Synlett.* 1992;1992(10):835-6.
24. Chinchilla R. In: Bäckvall JE, editor. *The chemistry of enamines.* New York: Wiley; 1994.
25. Whitesell JK, Whitesell MA. Alkylation of ketones and aldehydes via their nitrogen derivatives. *Synthesis.* 1983;1983(7):517-36.
26. Cook G. *Enamines: synthesis, structure, and reactions.* Boca Raton: CRC Press; 2017.
27. Zhang Y, Qi Y, Ulrich S, Barboiu M, Ramström O. Dynamic covalent polymers for biomedical applications. *Mater Chem Front.* 2020;4(2):489-506.
28. Frei P, Hevey R, Ernst B. Dynamic combinatorial chemistry: a new methodology comes of age. *Chem Eur J.* 2019;25(1):60-73.
29. Zhang Y, Zhang Y, Ramström O. Dynamic covalent kinetic resolution. *Catal Rev.* 2020;62(1):66-95.
30. Orrillo AG, Escalante AM, Martinez-Amezaga M, Cabezudo I, Furlan RL. Molecular networks in dynamic multilevel systems. *Chem Eur J.* 2019;25(5):1118-27.
31. Schaufelberger F, Timmer BJJ, Ramström O, Zhang W, Jin Y. *Dynamic Covalent Chemistry: Principles, Reactions, and Applications.* Weinheim: Wiley-VCH; 2017.
32. Miljanić OŠ. Small-molecule systems chemistry. *Chem.* 2017;2(4):502-24.
33. Kosikova T, Philp D. Exploring the emergence of complexity using synthetic replicators. *Chem Soc Rev.* 2017;46(23):7274-305.
34. Azcune I, Odriozola I. Aromatic disulfide crosslinks in polymer systems: Self-healing, reprocessability, recyclability and more. *Eur Polym J.* 2016;84:147-60.
35. Brachvogel RC, von Delius M. The dynamic covalent chemistry of esters, acetals and orthoesters. *Eur J Org Chem.* 2016;2016(22):3662-70.
36. Zhang Y, Barboiu M. Constitutional Dynamic Materials—Toward Natural Selection of Function. *Chem Rev.* 2016;116(3):809-34.
37. Lehn JM. Perspectives in chemistry—aspects of adaptive chemistry and materials. *Angew Chem Int Ed.* 2015;54(11):3276-89.
38. Herrmann A. Dynamic combinatorial/covalent chemistry: a tool to read, generate and modulate the bioactivity of compounds and compound mixtures. *Chem Soc Rev.* 2014;43(6):1899-933.
39. Hu L, Schaufelberger F, Timmer BJJ, Flos MA, Ramström O. *Kirk-Othmer Encyclopedia of Chemical Technology.* Hoboken: Wiley; 2014. p. 1-25.
40. Misuraca MC, Moulin E, Ruff Y, Giuseppone N. Supramolecular polymers: From structural control to functional materials. *New J Chem.* 2014;38(8):3336-49.
41. Wilson A, Gasparini G, Matile S. Functional systems with orthogonal photo- and redox inputs. *Chem Soc Rev.* 2014;43(5):1948-62.
42. Jin Y, Yu C, Denman RJ, Zhang W. Recent advances in dynamic covalent chemistry. *Chem Soc Rev.* 2013;42(16):6634-54.
43. Hunt RA, Otto S. Dynamic combinatorial libraries: Functional assembly in chemistry and biology. *Chem Commun.* 2011;47(3):847-58.
44. Gasparini G, Dal Molin M, Prins LJ. Dynamic covalent chemistry: From discrete molecules to functional supramolecular systems. *Eur J Org Chem.* 2010;2010(23):2429-40.
45. Kogan EA, Kuznetsov MT. Enamine compounds as potential bioactive agents: Recent advances. *J Med Chem.* 2019;62(10):4534-50.